




G BD4/1812



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ




**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 25 May 2004



The  
Patent  
Office

02MAY03 E804336-3, D00019  
P01/7700 0100-0310057.1

1/77

02MAY03 E804336-3, D00019  
P01/7700 0100-0310057.1  
The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

# Request for grant of a patent

See the notes on the back of this form. You can also  
get an explanatory leaflet from the Patent Office to  
help you fill in this form)

1. Your reference

P034018GB:BRC

2. Patent application number

(The Patent Office will fill in this part)

01 MAY 2003

0310057 5

3. Full name, address and postcode of the or of  
each applicant (underline all surnames)

VERNALIS RESEARCH LIMITED  
OAKDENE COURT  
613 READING ROAD  
WINNERSH  
WOKINGHAM  
RG41 5UA

Patents ADP number (if you know it)

07915184001

If the applicant is a corporate body, give the  
country/state of its incorporation

UNITED KINGDOM

4. Title of the invention

NEW USE FOR CHEMICAL COMPOUNDS

5. Name of your agent (if you have one)

Carpmaels & Ransford

"Address for service" in the United Kingdom  
to which all correspondence should be sent  
(including the postcode)

43 Bloomsbury Square  
London  
WC1A 2RA

Patents ADP number (if you know it)

83001

6. If you are declaring priority from one or more  
earlier patent applications, give the country  
and the date of filing of the or of each of these  
earlier applications and (if you know it) the or  
each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise  
derived from an earlier UK application,  
give the number and the filing date of  
the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right  
to grant of a patent required in support of  
this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an  
applicant, or
- c) any named applicant is a corporate body

Yes

See note (d))

**Patents Form 1/77**

9. Indicate the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	-
Description	28
Claim(s)	2
Abstract	-
Drawing(s)	-

*g*

10. If you are also filing any of the following, state how many against each item.

Priority documents	-
Translations of priority documents	-
Statement of inventorship and right to grant of a patent ( <i>Patents Form 7/77</i> )	-
Request for preliminary examination and search ( <i>Patents Form 9/77</i> )	-
Request for substantive examination ( <i>Patents Form 10/77</i> )	-
Any other documents ( <i>please specify</i> )	-

11. I/We request the grant of a patent on the basis of this application.

Signature

Date

*Cammaels & Ransford*  
Cammaels & Ransford

1st May 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

DR. BRUCE R. COCKERTON

020-7242 8692

**Warning**

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- For details of the fee and ways to pay please contact the Patent Office.*

NEW USE FOR CHEMICAL COMPOUNDS

The present invention relates primarily to the use of azetidine-1-carboxamides in the treatment of disorders mediated by the cannabinoid CB<sub>1</sub> receptor, particularly to the treatment of obesity and other eating disorders associated with excessive food intake.

It has been recognised that obesity is a disease process influenced by environmental factors in which the traditional weight loss methods of dieting and exercise need to be supplemented by therapeutic products (S. Parker, "Obesity: Trends and Treatments", Scrip Reports, PJB Publications Ltd, 1996).

Whether someone is classified as overweight or obese is generally determined on the basis of their body mass index (BMI) which is calculated by dividing body weight (kg) by height squared (m<sup>2</sup>). Thus, the units of BMI are kg/m<sup>2</sup> and it is possible to calculate the BMI range associated with minimum mortality in each decade of life. Overweight is defined as a BMI in the range 25-30 kg/m<sup>2</sup>, and obesity as a BMI greater than 30 kg/m<sup>2</sup>. There are problems with this definition in that it does not take into account the proportion of body mass that is muscle in relation to fat (adipose tissue). To account for this, obesity can also be defined on the basis of body fat content: greater than 25% and 30% in males and females, respectively.

As the BMI increases there is an increased risk of death from a variety of causes that is independent of other risk factors. The most common diseases with obesity are cardiovascular disease (particularly hypertension), diabetes (obesity aggravates the development of diabetes), gall bladder disease (particularly cancer) and diseases of reproduction. Research has shown that even a modest reduction in body weight can correspond to a significant reduction in the risk of developing coronary heart disease.

Compounds marketed as anti-obesity agents include Orlistat (Reductil®) and Sibutramine. Orlistat (a lipase inhibitor) inhibits fat absorption directly and tends to produce a high incidence of unpleasant (though relatively harmless) side-effects such as diarrhoea. Sibutramine (a mixed 5-HT/noradrenaline reuptake inhibitor) can increase blood pressure and heart rate in some patients. The serotonin releaser/reuptake inhibitors fenfluramine

(Pondimin®) and dexfenfluramine (Redux™) have been reported to decrease food intake and body weight over a prolonged period (greater than 6 months). However, both products were withdrawn after reports of preliminary evidence of heart valve abnormalities associated with their use. There is therefore a need for the development of a safer anti-  
5 obesity agent.

There now exists extensive pre-clinical and clinical data supporting the use of CB<sub>1</sub> receptor antagonists / inverse agonists for the treatment of obesity.

10 Preparations of marijuana (*Cannabis sativa*) have been used for over 5000 years for both medicinal and recreational purposes. The major psychoactive ingredient of marijuana has been identified as  $\Delta^9$ -tetrahydrocannabinol ( $\Delta^9$ -THC), one of a member of over 60 related cannabinoid compounds isolated from this plant. It has been demonstrated that  $\Delta^9$ -THC exerts its effects via agonist interaction with cannabinoid (CB) receptors. So far, two  
15 cannabinoid receptor subtypes have been characterised (CB<sub>1</sub> and CB<sub>2</sub>). The CB<sub>1</sub> receptor subtype is found predominantly in the central nervous system, and to a lesser extent in the peripheral nervous system and various peripheral organs. The CB<sub>2</sub> receptor subtype is found predominantly in lymphoid tissues and cells. To date, three endogenous agonists (endocannabinoids) have been identified which interact with both CB<sub>1</sub> and CB<sub>2</sub> receptors  
20 (anandamide, 2-arachidonyl glycerol and noladin ether).

Genetically obese rats and mice exhibit markedly elevated endocannabinoid levels in brain regions associated with ingestive behaviour (Di Marzo et al. 2001 Nature 410: 822 - 825). Furthermore, increased levels of endocannabinoids are observed upon the fasting of  
25 normal, lean animals (Kirkham et al., British Journal of Pharmacology, 2002, 136(4), 550-557). Exogenous application of endocannabinoids leads to the same physiological effects observed with  $\Delta^9$ -THC treatment, including appetite stimulation (Jamshida et al., British Journal of Pharmacology, 2001, 134: 1151-1154), analgesia, hypolocomotion, hypothermia, and catalepsy.

30

CB<sub>1</sub> (CB<sub>1</sub><sup>-/-</sup>) and CB<sub>2</sub> (CB<sub>2</sub><sup>-/-</sup>) receptor knockout mice have been used to elucidate the specific role of the two cannabinoid receptor subtypes. Furthermore, for ligands such as  $\Delta^9$ -THC which act as agonists at both receptors, these mice have allowed identification of

which receptor subtype is mediating specific physiological effects.  $CB_1^{-/-}$ , but not  $CB_2^{-/-}$ , mice are resistant to the behavioural effects of agonists such as  $\Delta^9$ -THC.  $CB_1^{-/-}$  animals have also been shown to be resistant to both the body weight gain associated with chronic high fat diet exposure, and the appetite-stimulating effects of acute food deprivation.

5

These findings suggest a clear role for both endogenous and exogenous cannabinoid receptor agonists in increasing food intake and body weight *via* selective activation of the  $CB_1$  receptor subtype.

- 10 The therapeutic potential for cannabinoid receptor ligands has been extensively reviewed (Exp. Opin. Ther. Pat. 1998, 8, 301-313; Exp. Opin. Ther. Pat. 2000, 10, 1529-1538; Trends in Pharm. Sci. 2000, 21, 218-224; Exp. Opin. Ther. Pat. 2002, 12(10), 1475-1489).

At least one compound (SR-141716A) characterised as a  $CB_1$  receptor antagonist / inverse

- 15 agonist is known to be in clinical trials for the treatment of obesity.

WO 00/15609, WO 01/64632, WO 01/64633 and WO 01/64634 disclose azetidine derivatives as  $CB_1$  receptor antagonists. WO 02/28346 discloses the association of an azetidine derivative as a  $CB_1$  receptor antagonist, and sibutramine, for the treatment of

20 obesity.

There remains a medical need for low molecular weight  $CB_1$  receptor antagonists / inverse agonists with pharmacokinetic and pharmacodynamic properties making them suitable for use as pharmaceutical agents. There also remains a medical need for new treatments of

- 25 disorders mediated by the  $CB_1$  receptor, particularly eating disorders, and particularly obesity. The object of the present invention is to provide such pharmaceutical agents and treatments.


It has now been found that certain azetidine-1-carboxamides show unexpected efficacy as

30

anti-obesity agents. These compounds were previously described in WO-A-99/37612 for the treatment of anxiety and epilepsy. These azetidine-1-carboxamides have been shown to selectively bind to the  $CB_1$  receptor subtype with high affinity. Such compounds have been shown to dose-dependently block the effects of an exogenously applied cannabinoid

receptor agonist (eg  $\Delta^9$ -THC) in mice. Furthermore, such compounds have been shown to reduce food intake and body weight gain in both rat and mouse models of feeding behaviour.

- 5 According to the present invention, there is provided use of a compound of formula (I)



wherein:

R<sup>1</sup> is aryl;

R<sup>2</sup> is H, alkyl or aryl; and

- 10 R<sup>3</sup> is hydrogen or alkyl;

or a pharmaceutically acceptable salt or prodrug thereof, in the manufacture of a medicament for the treatment of a disorder mediated by CB<sub>1</sub> receptors.

- The active compounds of formula (I) are antagonists and/or inverse agonists at the  
 15 cannabinoid-1 (CB<sub>1</sub>) receptor and are useful for the treatment, prevention and suppression of diseases mediated by the CB<sub>1</sub> receptor. The invention is concerned with the use of these compounds to selectively antagonise the CB<sub>1</sub> receptor and, as such, in the treatment of obesity and other disorders.

- 20 Reference in the present specification to an "alkyl" group means a branched or unbranched, cyclic or acyclic, saturated or unsaturated (e.g. alkenyl (including allyl) or alkynyl (including propargyl)) hydrocarbyl radical. Where cyclic or acyclic the alkyl group is preferably C<sub>1</sub> to C<sub>12</sub>, more preferably C<sub>1</sub> to C<sub>8</sub> (such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, sec-butyl, pentyl, isopentyl, hexyl, heptyl, octyl). It will be appreciated therefore that  
 25 the term "alkyl" as used herein includes alkyl (branched or unbranched), alkenyl (branched or unbranched), alkynyl (branched or unbranched), cycloalkyl, cycloalkenyl and cycloalkynyl. A cyclic alkyl group may also be a mono-bridged or multi-bridged cyclic alkyl group. In a preferred embodiment, a cyclic alkyl group is preferably C<sub>3</sub> to C<sub>12</sub>, more preferably C<sub>5</sub> to C<sub>8</sub> and an acyclic alkyl group is preferably C<sub>1</sub> to C<sub>10</sub>, more preferably C<sub>1</sub> to C<sub>6</sub>, more preferably

methyl, ethyl, propyl (n-propyl or isopropyl), butyl (n-butyl, isobutyl, tertiarybutyl or sec-butyl) or pentyl (including n-pentyl and iso-pentyl), more preferably methyl.

- As used herein, the term "lower alkyl" means a branched or unbranched, cyclic or acyclic, saturated or unsaturated (e.g. alkenyl or alkynyl) hydrocarbyl radical wherein said cyclic lower alkyl group is C<sub>5</sub>, C<sub>6</sub> or C<sub>7</sub>, and wherein said acyclic lower alkyl group is C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub> or C<sub>4</sub>. It will be appreciated therefore that the term "lower alkyl" as used herein includes lower alkyl (branched or unbranched), lower alkenyl (branched or unbranched), lower alkynyl (branched or unbranched), cycloloweralkyl, cycloloweralkenyl and cycloloweralkynyl.
- 10 Preferably, a lower alkyl group is preferably selected from methyl, ethyl, propyl (n-propyl or isopropyl) or butyl (n-butyl, isobutyl, sec-butyl or tertiary-butyl), preferably methyl.

Reference in the present specification to an "aryl" group means a mono or bicyclic aromatic group, such as phenyl or naphthyl, and preferably a mono-cyclic aromatic group.

- 15 Reference in the present specification to a "heteroaryl" group means an aromatic group containing one or more heteroatoms, preferably 1, 2 or 3 heteroatoms, preferably 1 or 2 heteroatoms. Preferably the heteroatoms are selected from O, S and N, preferably from O and N. Preferably the heteroaryl group comprises 5 or 6-membered ring systems. The heteroaryl
- 20 group is preferably a monocyclic or bicyclic ring system, preferably monocyclic. Examples include thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl and isobenzofuryl.

- 25 Reference in the present specification to a non-aromatic heterocyclic group is to a saturated or partially unsaturated 4, 5, 6 or 7-membered ring containing 1, 2 or 3 heteroatoms selected from N, O and S, preferably 1 or 2 heteroatoms, preferably selected from N and O. Examples include piperidinyl, morpholinyl, piperazinyl and pyrrolidinyl.

- The alkyl and aryl groups may be substituted or unsubstituted. In one embodiment, only the
- 30 alkyl and aryl groups defined herein as R<sup>1</sup> to R<sup>3</sup> and R<sup>9</sup> to R<sup>13</sup> may be substituted. Where substituted, there will generally be 1 to 3 substituents present, preferably 1 or 2 substituents. Substituents may include:



carbon containing groups such as

alkyl

aryl, arylalkyl (e.g. substituted and unsubstituted phenyl, substituted and unsubstituted benzyl);

5 halogen atoms and halogen containing groups such as

haloalkyl (e.g. trifluoromethyl);

oxygen containing groups such as

alcohols (e.g. hydroxy, hydroxyalkyl, (aryl)(hydroxy)alkyl),

ethers (e.g. alkoxy, alkoxyalkyl, aryloxyalkyl),

10 aldehydes (e.g. carboxaldehyde),

ketones (e.g. alkylcarbonyl, alkylcarbonylalkyl, arylcarbonyl, arylalkylcarbonyl, arylcarbonylalkyl),

acids (e.g. carboxy, carboxyalkyl),

acid derivatives such as esters

15 (e.g. alkoxycarbonyl, alkoxycarbonylalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl)

and amides

(e.g. aminocarbonyl, mono- or dialkylaminocarbonyl, aminocarbonylalkyl, mono- or dialkylaminocarbonylalkyl, arylaminocarbonyl);

20

nitrogen containing groups such as

amines (e.g. amino, mono- or dialkylamino, aminoalkyl, mono- or dialkylaminoalkyl),

azides,

25 nitriles (e.g. cyano, cyanoalkyl),

nitro;

sulphur containing groups such as

thiols, thioethers, sulphoxides and sulphones

(e.g. alkylthio, alkylsulfinyl, alkylsulfonyl, alkylthioalkyl, alkylsulfinylalkyl,

30

alkylsulfonylalkyl, arylthio, arylsulfinyl, arylsulfonyl, arylthioalkyl, arylsulfinylalkyl, arylsulfonylalkyl);

and heterocyclic groups containing one or more, preferably one, heteroatom,

(e.g. thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazoliny, pyrazolidinyl, tetrahydrofuranyl, pyranyl, pyronyl, pyridyl, pyrazinyl, pyridazinyl, piperidyl, piperazinyl, morpholinyl, thionaphthyl, benzofuranyl, isobenzofuryl, indolyl, oxyindolyl, isoindolyl, indazolyl, indolinyl, 7-azaindolyl, isoindazolyl, benzopyranyl, coumarinyl, isocoumarinyl, quinolyl, isoquinolyl, naphthridinyl, cinnolinyl, quinazolinyl, pyridopyridyl, benzoxazinyl, quinoxadinyl, chromenyl, chromanyl, isochromanyl and carbolinyl).

Where an aryl group is phenyl, the phenyl may be substituted by adjacent substituents forming a 5 or 6 membered saturated ring optionally containing 1 or 2 heteroatoms, preferably selected from N, O and S, preferably from N and O. Where the saturated ring contains 2 nitrogen atoms, the ring is preferably a 6-membered ring. Where the saturated ring contains 2 oxygen atoms, the ring may be a 5- or 6-membered ring. Examples include 2,3-dihydrobenzo[b]furan-7-yl, 2,3-dihydrobenzo[b]thiophen-6-yl, 1,2,3,4-tetrahydronaphthalen-5-yl, 2,3-dihydrobenzo[1,4]dioxin-6-yl and 1,2,3,4-tetrahydroisoquinolin-8-yl.

Preferred substituents include alkyl (including haloalkyl), alkoxy (including haloalkoxy), aryl, nitrile or halo. Preferred halogen-containing groups include trifluoromethyl.

As used herein, the term "alkoxy" means alkyl-O- and "alkoyl" means alkyl-CO-.

As used herein, the term "halogen" means a fluorine, chlorine, bromine or iodine radical, preferably a fluorine or chlorine radical.

The compounds of formula (I) may exist in a number of diastereomeric and/or enantiomeric forms. Unless otherwise stated, reference in the present specification to "a compound of formula (I)" is a reference to all stereoisomeric forms of the compound and includes a

reference to the unseparated stereoisomers in a mixture, racemic or non-racemic, and to each stereoisomer in its pure form.

- In the compounds of formula (I), preferably  $R^1$  is substituted or unsubstituted phenyl or naphthyl (preferably phenyl), more preferably  $R^1$  is a substituted phenyl or naphthyl (preferably phenyl), more preferably  $R^1$  is a phenyl or naphthyl (preferably phenyl), having 1 to 3 substituents and most preferably  $R^1$  is a phenyl or naphthyl (preferably phenyl), having 1 or 2 substituents. Preferred substituents include alkyl, halo, halogen-containing groups such as haloalkyl (particularly halomethyl, such as trifluoromethyl), thioalkyl, alkoxy, alkylsulfonyl, and mono- or di-alkylaminocarbonyl. Particularly preferred substituents are alkyl, halo and halogen-containing groups such as haloalkyl (particularly halomethyl, such as trifluoromethyl); more preferably halo and halogen-containing groups such as haloalkyl (particularly halomethyl, such as trifluoromethyl).
- 15 In one embodiment of the invention,  $R^2$  is aryl, preferably substituted or unsubstituted phenyl, more preferably substituted phenyl, more preferably phenyl having 1 to 3 substituents and most preferably phenyl having 1 or 2 substituents. Preferred substituents include alkyl, halo, halogen-containing groups such as haloalkyl (particularly halomethyl, such as trifluoromethyl), thioalkyl, alkoxy, alkylsulfonyl, and mono- or di-alkylaminocarbonyl.
- 20 Particularly preferred substituents are alkyl, halo and halogen-containing groups such as haloalkyl (particularly halomethyl, such as trifluoromethyl); more preferably halo and halogen-containing groups such as haloalkyl (particularly halomethyl, such as trifluoromethyl).
- 25 In an alternative embodiment,  $R^2$  is H or alkyl (cyclic or acyclic).

- In a preferred embodiment,  $R^3$  is alkyl, and preferably selected from alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl or unsubstituted saturated cyclic or acyclic hydrocarbyl. Preferably  $R^3$  is acyclic hydrocarbyl, preferably lower alkyl, and in one embodiment is substituted. One or two substituent groups may be present, preferably one substituent group. Preferred substituents are those referred to hereinabove, particularly hydroxy, alkoxy, thioalkyl, amino, mono- and dialkyl amino, alkoxycarbonyl, aryl (preferably phenyl), and heterocyclic groups including both heteroaryl and non-aromatic heterocyclic groups. Where  $R^3$  is an acyclic alkyl
- 30

group, it may be substituted by a cyclic alkyl group; and where  $R^3$  is a cyclic alkyl group it may be substituted by an acyclic alkyl group. Where the substituent group is heteroaryl, the heteroaryl preferably is a 5- or 6-membered ring containing one or more N, O or S atoms, and preferred groups include thiophenyl, furanyl, isoxazolyl, thiazolyl and benzothiophenyl. Other preferred substituent groups include dihydrobenzofuranyl, dihydrobenzodioxinyl, tetrahydrofuranyl, pyrrolidinyl, oxopyrrolindyl and benzodioxolyl.

In one embodiment,  $R^3$  is selected from :



wherein  $n$  is 0 or 1;

$m$  is 0, 1, 2 or 3;

$R^9$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are selected from hydrogen, alkyl (preferably lower alkyl), hydroxy, alkoxy (preferably lower alkoxy), thioalkyl (preferably thio lower alkyl), amino, mono- and di-alkyl amino (preferably lower alkyl amino), alkoxycarbonyl (preferably lower alkoxy carbonyl) and  $R^{13}$ ;

wherein  $R^{13}$  is selected from aryl, heteroaryl and non-aromatic heterocyclic optionally substituted by one or more (preferably 1 or 2, preferably 1) groups preferably selected from alkyl (preferably lower alkyl, preferably methyl), halogen (preferably fluoro, chloro and bromo), alkoxy (preferably lower alkoxy, preferably methoxy), oxo, aryl, heteroaryl and non-aromatic heterocycle.

Preferably,  $m$  is 0 or 1 or 2, preferably 0 or 1, and preferably 0.

Preferably,  $n$  is 0.

In one embodiment, at least one and more preferably two of  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are selected from hydrogen. In a further embodiment, at least one and more preferably at least two of  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  are selected from methyl.

In a further embodiment,  $R^3$  is selected from cyclic alkyl, including cyclopentyl, cyclohexyl, norbornanyl and adamantyl, preferably cyclopentyl and cyclohexyl.

Preferred R<sup>3</sup> groups are tertiary butyl, sec-butyl, isobutyl, isopropyl, n-propyl and ethyl, particularly tertiary butyl, isobutyl, sec-butyl and isopropyl, and particularly tertiary butyl.

Compounds of formula (I) include:

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
4-Cl-C <sub>6</sub> H <sub>4</sub>	4-Cl-C <sub>6</sub> H <sub>4</sub>	Allyl
4-Cl-C <sub>6</sub> H <sub>4</sub>	4-Cl-C <sub>6</sub> H <sub>4</sub>	2-Hydroxypropyl

5

According to a further aspect of the present invention there is provided a method of treatment of a disorder mediated by CB<sub>1</sub> receptors comprising administration to a subject in need of such treatment an effective dose of the compound of formula (I), or a  
10 pharmaceutically acceptable salt or prodrug thereof.

The diseases and disorders to which the present invention is directed are:  
psychosis, schizophrenia, cognitive disorders, attention deficit disorder, gastrointestinal disorders (such as dysfunction of gastrointestinal motility or diarrhoea), smoking cessation,  
15 obesity and other eating disorders associated with excessive food intake (including bulimia and compulsive eating disorder) and associated health complications including non-insulin dependant diabetes mellitus. The present invention is particularly directed to obesity and other eating disorders associated with excessive food intake and associated health complications including non-insulin dependant diabetes mellitus, and particularly to  
20 obesity and other eating disorders associated with excessive food intake, and especially to obesity.

25

In an alternative embodiment, the present invention is directed to smoking cessation and the facilitation thereof.

In a further alternative embodiment, the present invention is directed to gastrointestinal disorders (such as dysfunction of gastrointestinal motility or diarrhoea).

The present invention may be employed in respect of a human or animal subject, more  
30 preferably a mammal, more preferably a human subject.

As used herein, the term "treatment" as used herein includes prophylactic treatment.

As used herein, the term "prodrug" means any pharmaceutically acceptable prodrug of the compound of formula (I). For example, the compound of formula (I) may be prepared in a prodrug form wherein a free -OH group is derivatised (for example, via an ester, amide or phosphate bond) with a suitable group (the group may contain, for example, an alkyl, aryl, phosphate, sugar, amine, glycol, sulfonate or acid function) which is suitably labile so as it will be removed / cleaved (eg. by hydrolysis) to reveal the compound of formula (I) sometime after administration or when exposed to the desired biological environment.


As used herein, the term "pharmaceutically acceptable salt" means any pharmaceutically acceptable salt of the compound of formula (I). Salts may be prepared from pharmaceutically acceptable non-toxic acids and bases including inorganic and organic acids and bases. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethenesulfonic, dichloroacetic, fumaric, gluconic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, oxalic, p-toluenesulfonic and the like. Particularly preferred are hydrochloric, hydrobromic, phosphoric, sulfuric and methanesulfonic acids, and most particularly preferred is the methanesulfonate salt. Acceptable base salts include alkali metal (e.g. sodium, potassium), alkaline earth metal (e.g. calcium, magnesium) and aluminium salts.

The compound of formula (I) may be used in combination with one or more additional drugs useful in the treatment of the disorders mentioned above, the components being in the same formulation or in separate formulations for administration simultaneously or sequentially.

Compounds of formula (I) may be prepared according to Reaction Scheme 1 (where P is a nitrogen protecting group).  $R^1$ ,  $R^2$ , and  $R^3$  are as previously defined. The ether (IV) may be formed by reaction of the azetidinol (II) either with an arylalkanol (III,  $X = OH$ ) and diethylazo dicarboxylate and triphenyl phosphine or with an arylalkyl chloride, bromide, iodide, mesylate or tosylate (III,  $X = Cl, Br, I, mesylate, tosylate$ ) and a strong base such as

sodium hydride. Formation of the azetidine (V) may be achieved by reaction of (IV) with a suitable nitrogen deprotection agent. For example, if P is a diphenylmethyl group, then deprotection may be carried out by treatment with 1-chloroethyl chloroformate followed by methanol. The urea (I) is formed by reaction of azetidine (V) with an N-alkylisocyanate or an N-alkylcarbamoyl chloride and a base such as triethylamine or potassium carbonate. Alternatively, the urea may be prepared directly from the azetidine (IV) without isolation of an intermediate such as the secondary amine (V). For example, when P is a diphenylmethyl group, azetidine (IV) may be treated with phosgene followed by amine  $R^3NH_2$  to give urea (I) directly.

10

**Reaction Scheme 1**


Compounds of formula (I) where  $R^1$  and  $R^2$  are aryl may be prepared according to Reaction Scheme 2 (where P is a nitrogen protecting group).  $R^1$ ,  $R^2$ , and  $R^3$  are as previously described. The ether (IV) may be formed by reaction of the azetidinol (II) with a benzhydrol (III,  $X = OH$ ) with removal of water (for example azeotropic removal of water under standard Dean-Stark conditions). The ether (IV) may also be formed by reaction of the azetidinol (II) with a benzhydryl group substituted with a suitable leaving-group (III,  $X = Cl, Br, I, \text{mesylate, tosylate}$ ) and a strong base such as sodium hydride. Formation of the azetidine (V) may be achieved by reaction of (IV) with a suitable nitrogen deprotection agent. For example, if P is a benzhydryl group, then deprotection may be carried out by treatment with 1-chloroethyl chloroformate followed by treatment with methanol. The deprotected azetidine (V) can be

20

isolated directly as the hydrochloride salt or, upon basification, as the free-base. The urea (I) can be formed by reaction of azetidine (V) with an N-alkyl isocyanate, or an N-alkyl carbamoyl chloride and a base such as triethylamine or potassium carbonate. Alternatively, the urea may be prepared directly from the protected azetidine (IV) without isolation of the intermediate azetidine (V). For example, when P is a benzhydryl group, azetidine (IV) may be treated with phosgene followed by an amine,  $R^3NH_2$ , to give urea (I) directly. Azetidine (V) may also be converted to the corresponding carbamoyl chloride (VI) by treatment with, for example, triphosgene. This intermediate carbamoyl chloride (VI) may be reacted with an amine,  $R^3NH_2$ , to give the urea (I).

10

### Reaction Scheme 2



15 The invention further provides a pharmaceutical composition comprising an effective amount of the compound of formula (I) in combination with a pharmaceutically acceptable carrier or excipient and a method of making such a composition comprising combining an effective amount of the compound of formula (I) with a pharmaceutically acceptable carrier or excipient.

20



To further increase efficacy, the composition may contain components such as dextrans or cyclodextrins or ether derivatives thereof, which aid stability and dispersion, and decrease metabolism of the active ingredient.

- 5 For compositions in which the pharmaceutically acceptable carrier comprises a cyclodextrin or an ether derivative thereof, the active ingredient is intimately mixed with an aqueous solution of the cyclodextrin or ether derivative thereof, with optional addition of further pharmaceutically acceptable ingredients before, during or after said mixing. The thus obtained solution is optionally lyophilized, and the lyophilized residue is optionally  
10 reconstituted with water.

In an embodiment of the present invention, the composition further comprises a buffer system, an isotonicizing agent and water.

- 15 Compounds of formula (I) may be administered in a form suitable for oral use, for example a tablet, capsule, aqueous or oily solution, suspension or emulsion; for topical use including transmucosal and transdermal use, for example a cream, ointment, gel, aqueous or oil solution or suspension, salve, patch or plaster; for nasal use, for example a snuff, nasal spray or nasal drops; for vaginal or rectal use, for example a suppository; for administration by inhalation,  
20 for example a finely divided powder or a liquid aerosol; for sub-lingual or buccal use, for example a tablet or capsule; or for parenteral use (including intravenous, subcutaneous, intramuscular, intravascular or infusion), for example a sterile aqueous or oil solution or suspension. In general the above compositions may be prepared in a conventional manner using conventional excipients, using standard techniques well known to those skilled in the art  
25 of pharmacy. Preferably, the compound is administered orally.

For oral administration, the compounds of formula (I) will generally be provided in the form of tablets or capsules or as an aqueous solution or suspension.

- 30 Tablets for oral use may include the active ingredient mixed with pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavouring agents, colouring agents and preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and calcium phosphate, and lactose,

while corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin, while the lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate, to delay absorption in the gastrointestinal tract.

5

Capsules for oral use include hard gelatin capsules in which the active ingredient is mixed with a solid diluent, and soft gelatin capsules wherein the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin or olive oil.

- 10 For intramuscular, intraperitoneal, subcutaneous and intravenous use, the compounds of formula (I) will generally be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and isotonicity. Suitable aqueous vehicles include Ringer's solution and isotonic sodium chloride. Aqueous suspensions may include suspending agents such as cellulose derivatives, sodium alginate, polyvinyl-pyrrolidone and gum tragacanth, and a  
15 wetting agent such as lecithin. Suitable preservatives for aqueous suspensions include ethyl and n-propyl p-hydroxybenzoate.

It will be appreciated that the dosage levels used may vary over quite a wide range depending upon the compound used, the severity of the symptoms exhibited by the patient and the  
20 patient's body weight.

The invention will now be described in detail with reference to the following pharmacological examples. It will be appreciated that the examples are intended to illustrate and not to limit the scope of the present invention.

25

## EXAMPLES

### Synthetic Examples

- 30 Preparation of 1-(Diphenylmethyl)-3-azetidinol

This compound was prepared according to the method of Anderson and Lok (*J. Org. Chem.*, 1972, 37, 3953, the disclosure of which is incorporated herein by reference), m.p. 111-112 °C (lit. m.p. 113 °C).

**Preparation of 3-(4-Chlorobenzoyloxy)-1-(diphenylmethyl) azetidine (1)**

- 5 A solution of 1-diphenylmethyl-3-azetidinol (25 mmol) in DMF (100 mL) was added at 0 °C to a suspension of NaH (60% disp.in oil, 30 mmol) in DMF (50 mL). The reaction mixture was stirred at room temperature for 1h, then 4-chlorobenzylchloride (25 mmol) was added dropwise at 0 °C and the reaction mixture stirred at room temperature for 3 h. The reaction was quenched with water and extracted with ethyl acetate (3 x 50 mL), the extracts were
- 10 washed with water and brine, dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. The residue was purified by chromatography [SiO<sub>2</sub>; hexane-ethyl acetate (9:1)] to yield the product as a yellow oil (7.3 g, 80%). The material was used in the next step without further purification.

**Example 1. 3-(4-Chlorobenzoyloxy)-N-(2-propenyl)azetidine-1-carboxamide (2)**

- Phosgene solution (1.75-M in toluene, 24 mmol) was added at 0°C to a solution of compound
- 15 (1) (20 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL). The reaction mixture was stirred at room temperature for 90 min, concentrated *in vacuo*, then redissolved in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and treated with allylamine (42 mmol) at 0°C. The reaction was stirred for 4 h at room temperature, then water (40 mL) was added and the layers were separated. The aqueous layer was extracted with further CH<sub>2</sub>Cl<sub>2</sub> (2 x 40 mL). The organic layers were washed with dilute HCl (20 mmol)
- 20 and brine, dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. The residue was triturated using diethyl ether to give the product (2) as a crystalline solid (3.5 g, 60%), m.p. 110-111 °C. Found: C, 59.84; H, 6.11; N, 9.98. C<sub>14</sub>H<sub>17</sub>ClN<sub>2</sub>O<sub>2</sub> requires: C, 59.89; H, 9.6.10; N, 9.97%.

**Preparation of 3-(3,4-Dichlorobenzoyloxy)-1-(diphenylmethyl) azetidine (3)**

- This material was prepared from 1-diphenylmethyl-3-azetidinol (6.0 g) and alpha,3,4-
- 25 trichlorotoluene using the procedure described for compound (1) (yield 92%).

**Example 2. 3-(3,4-Dichlorobenzoyloxy)-N-(2-propenyl)azetidine-1-carboxamide (4)**

This material was prepared from compound (3) (9.2 g) using the procedure described for compound (2) (yield 75%), m.p. 88-89 °C. Found: C, 53.43; H, 5.18; N, 8.85,  $C_{14}H_{16}Cl_2N_2O_2$  requires C, 53.35; H, 5.12; N, 8.88%.

**Preparation of 3-(3-(Trifluoromethyl)benzyloxy)-1-(diphenylmethyl)azetidine (5)**

- 5 This material was prepared from 1-diphenylmethyl-3-azetidinol (5 g) and  $\alpha'$ -bromo- $\alpha,\alpha,\alpha$ -trifluoro-*m*-xylene using the procedure described for compound (1) (yield 91%).

**Example 3. 3-(3-(Trifluoromethyl)benzyloxy)-N-(2-propenyl)azetidine-1-carboxamide (6)**

- 10 This material was prepared from compound (5) (7.5 g) using the procedure described for compound (1) (yield 64%), m.p. 108°C. Found: C, 57.29; H, 5.44; N, 8.87,  $C_{15}H_{17}F_3N_2O_2$  requires C, 57.32; H, 5.45; N, 8.91%.

**Preparation of 3-(4-(Trifluoromethyl)benzyloxy)-1-(diphenylmethyl)azetidine (7)**

- 15 This material was prepared from 1-diphenylmethyl-3-azetidinol (6.0 g) and  $\alpha'$ -bromo- $\alpha,\alpha,\alpha$ -trifluoro-*p*-xylene using the procedure described for compound (1) (yield 77%).

**Example 4. 3-(4-(Trifluoromethyl)benzyloxy)-N-(2-propenyl)azetidine-1-carboxamide (8)**

- 20 This material was prepared from compound (7) (7.7 g) using the procedure described for compound (2) (yield 72%), m.p. 120 °C. Found: C, 57.27; H, 5.45; N, 8.86.  $C_{15}H_{17}F_3N_2O_2$  requires C, 57.32; H, 5.45; N, 8.91%.

**Preparation of 3-(4-Fluorobenzyloxy)-1-(diphenylmethyl) azetidine (9)**

- 25 This material was prepared from 1-diphenylmethyl-3-azetidinol (6.0 g) and 4-fluorobenzyl bromide using the procedure described for compound (1) (yield 83%).

**Example 5. 3-(4-Fluorobenzyloxy)-N-(2-propenyl)azetidine-1-carboxamide (10)**

This material was prepared from compound (9) using the procedure described for compound (2), m.p. 97-99 °C. Found: C, 63.57; H, 6.59; N, 10.66.  $C_{14}H_{17}ClN_2O_2$  requires C, 63.62; H, 6.48; N, 10.59.

**5 Preparation of 3-(bis-(4-chlorophenyl)methoxy-1-diphenylmethyl)azetidine (11)**

A solution of 4,4'-dichlorobenzhydrol (25 mmol), *p*-toluenesulfonic acid (18.4 mmol) and 1-(diphenylmethyl)-3-azetidinol (8.4 mmol) in benzene (100 mL) was heated under reflux in a Dean-Stark apparatus for 3h. The solution was cooled, washed with sodium hydrogen carbonate (saturated aqueous solution, 100 mL), dried ( $MgSO_4$ ) and concentrated *in vacuo*.

- 10 The residue was purified by chromatography [ $SiO_2$ ; hexane-diethyl ether (5:1)] to yield the product (11) as a thick oil that crystallized on standing (2.4g, 62%).

**Example 6. 3-(Bis(4-chlorophenyl)methoxy)-N-(2-propenyl)azetidine-1-carboxamide (12)**

- 15 This material was prepared from compound (11) using the procedure described for compound (2) (yield 17%) as a crystalline solid. Found: C, 56.38; H, 5.10; N, 6.51.  $C_{20}H_{20}Cl_2N_2O_2 \cdot 2H_2O$  requires: C, 56.21; H, 5.66; N, 6.56%.

**Example 7. Preparation of (R)-3-(Bis(4-chlorophenyl)methoxy)-N-(2-**

**20 hydroxypropyl)azetidine-1-carboxamide (13)**

This material was prepared from compound (11) and (*R*)-(-)-1-amino-2-propanol using the procedure described for compound (2) (yield 57%) as a crystalline solid. Found: C, 58.74; H, 5.42; N, 6.84.  $C_{20}H_{22}Cl_2N_2O_3$  requires: C, 58.69; H, 5.42; N, 6.84%.

**25 Example 8. 3-(3-Trifluoromethyl)benzyloxy-N-azetidine-1-carboxamide (14)**

To a solution of 3-(3-trifluoromethyl)benzyloxy-1-(diphenylmethyl)azetidine (5) (5.3 mmol) in dichloromethane (15 mL) at 0°C, was added a solution of phosgene (1.75M in toluene, 6.4 mmol). The reaction mixture was stirred at room temperature for 2h, concentrated *in vacuo*, then redissolved in THF (15 mL) and treated with ammonium hydroxide (5 mL), added in one  
30 portion, at 0°C. The reaction was stirred vigorously for 15h at room temperature, then water (50 mL) and ethyl acetate (40 mL) were added and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 40 mL), dried ( $MgSO_4$ ) and concentrated *in vacuo*.

The residue was triturated using ethyl acetate (10 mL) to yield (14) as a solid (0.91 g, 63%), mp. 167 °C (ethyl acetate).

Found: C, 52.44; H, 4.72; N, 10.23.  $C_{14}H_{17}ClN_2O_2$  requires: C, 52.56; H, 4.78; N, 10.21.

5 **Preparation of 3-(1-(3-trifluoromethylphenyl)ethyloxy)-1-(diphenylmethyl)azetidine (15)**

To a solution of  $\alpha$ -methyl-3-trifluoromethylbenzyl alcohol (53 mmol), diisopropylethyl amine (105 mmol) in dichloromethane (150 mL) under nitrogen and cooled to 0 °C, was added methane sulfonyl chloride (63.1 mmol) dropwise over 10 min. The reaction was stirred for 15h. Water (200 mL) was added and the resulting mixture stirred for 10min, poured into  
10 potassium carbonate (10% wt/wt aqueous solution, 200 mL) and extracted with dichloromethane (3x150 mL). Combined organic extracts were washed with brine (50 mL) once and then dried ( $Na_2SO_4$ ), filtered and concentrated *in vacuo*. The residue was dissolved in ethyl ether and washed through a pad of silica, eluting with more ether. The filtrate was concentrated *in vacuo*. This material was used directly, as shown below.

15

A solution of 1-diphenylmethyl-3-azetidinol (42 mmol) in dimethyl formamide (20 mL) was added via pipette, to a suspension of NaH (60% disp.in oil, 50 mmol) in dimethyl formamide (80 mL) at 0°C. The reaction mixture was stirred at room temperature for 15 min, the crude material from above (assumed 53 mmol) was added dropwise as a solution in dimethyl  
20 formamide (30 mL) at 0°C and the reaction mixture stirred at room temperature for 2 h. The reaction was poured into water (200 mL) and extracted with ethyl acetate (3 x 50 mL), the extracts were washed with water (200 mL) and brine (50 mL), dried ( $MgSO_4$ ) and concentrated *in vacuo*. The residue was purified by chromatography ( $SiO_2$ ; hexane/ethyl acetate 9/1) to yield 3-(1-(3-trifluoromethylphenyl)ethyloxy)-1-(diphenylmethyl)azetidine  
25 (15) as a yellow oil (11.2g, yield 65%). The material was used in the next step without further purification.

**Example 9. 3-(1-(3-Trifluoromethylphenyl)ethyloxy)-azetidine-1-carboxamide (16)**

This material was prepared from compound (15) using the procedure described for compound  
30 (14) (yield 62%) as a crystalline solid, mp. 130.5-131.5°C (diisopropyl ether).









Found: C, 54.24; H, 5.26; N, 9.69.  $C_{14}H_{17}ClN_2O_2$  requires: C, 54.17; H, 5.24.; N, 9.71.

The individual enantiomers of Example 9 are prepared using the same overall synthetic method as described for compound 16, but using the chiral alcohols. The *R*-enantiomer of Example 9 was prepared from the appropriate chiral 1-(3-trifluoromethyl)phenyl ethyl alcohol. The chiral alcohols may be prepared from 3'-trifluoromethyl-acetophenone by  
5 stereoselective reduction, for example using borane and a suitable chiral auxiliary or chiral catalyst (see Corey, EJ; Bakshi, RK; Shibata S. *J. Amer. Chem. Soc.*, **1987**, *109*, 5551-5553 or Pickard, ST and Smith, HE. *J. Amer. Chem. Soc.*, **1990**, *112*, 5741-5747).









**Examples 10 to 43 – See Table 1**









10 These products were prepared using the procedure described for compound (2).









Table 1



Exempleno	Compound No.	Structure	Formula	MWt	mp	Cfound	Hfound	Nfound	Cexp	Hexp	Nexp	Note
10	17		C <sub>15</sub> H <sub>17</sub> N <sub>3</sub> O <sub>2</sub>	271.32	95-96	66.69	6.29	15.32	66.40	6.32	15.48	
11	18		C <sub>20</sub> H <sub>22</sub> N <sub>2</sub> O <sub>2</sub>	322.41	160.0	74.52	6.87	8.61	74.51	6.88	8.68	
12	19		C <sub>18</sub> H <sub>20</sub> N <sub>2</sub> O <sub>2</sub>	296.37	141-142	72.96	6.77	9.65	72.95	6.80	9.45	
13	20		C <sub>14</sub> H <sub>18</sub> O <sub>2</sub> N <sub>2</sub> O <sub>2</sub>	317.22	89-90	53.00	5.74	8.73	53.01	5.72	8.83	
14	21		C <sub>14</sub> H <sub>17</sub> ClN <sub>2</sub> O <sub>2</sub>	280.76	67-68	59.94	6.12	9.95	59.89	6.10	9.97	
15	22		C <sub>14</sub> H <sub>17</sub> FN <sub>2</sub> O <sub>2</sub>	264.30	59-60	63.55	6.55	10.59	63.62	6.48	10.59	
16	23		C <sub>15</sub> H <sub>19</sub> F <sub>3</sub> N <sub>2</sub> O <sub>2</sub>	316.33	128-129	56.92	6.09	8.83	56.96	6.05	8.85	
17	24		C <sub>15</sub> H <sub>18</sub> F <sub>3</sub> N <sub>2</sub> O <sub>2</sub>	316.33	62-63	56.89	6.21	8.82	56.96	6.05	8.85	



Exempleno	Compound No.	Structure	Formula	MWt	mp	Cfound	Hfound	Nfound	Cexp	Hexp	Nexp	Note
18	25		C15H19F3N2O3	332.33	67-68	54.25	5.81	8.42	54.21	5.76	8.43	
19	26		C15H19F3N2O3	332.33	67-68	54.21	5.87	8.41	54.21	5.76	8.43	
20	27		C15H19F3N2O3	332.33	97-98	54.09	5.76	8.39	54.21	5.76	8.43	
21	28		C15H19F3N2O3	332.33	97-98	54.39	5.82	8.44	54.21	5.76	8.43	
22	29		C14H18Cl2N2O3	333.22	88-89	50.46	5.34	8.39	50.46	5.44	8.40	
23	30		C14H18Cl2N2O3	333.22	88-89	50.49	5.36	8.61	50.46	5.44	8.40	
24	31		C14H19ClN2O3	298.77	85-86	56.27	6.40	9.35	56.28	6.41	9.37	
25	32		C15H15F3N2O2	312.29	90-91	57.73	4.94	8.91	57.69	4.84	8.97	

Exempleno	Compound No.	Structure	Formula	MWt	mp	C <sub>1</sub> ound	H <sub>1</sub> ound	N <sub>1</sub> ound	C <sub>exp</sub>	H <sub>exp</sub>	N <sub>exp</sub>	Note
26	33		C <sub>14</sub> H <sub>18</sub> N <sub>2</sub> O <sub>2</sub>	246.31	76-77	68.29	7.35	11.37	68.27	7.37	11.37	
27	34		C <sub>14</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>	282.32	73-74	59.49	6.87	9.93	59.56	6.78	9.92	
28	35		C <sub>15</sub> H <sub>17</sub> F <sub>3</sub> N <sub>2</sub> O <sub>2</sub>	314.31	63.0	57.34	5.47	8.92	57.32	5.45	8.91	
29	36		C <sub>14</sub> H <sub>16</sub> F <sub>2</sub> N <sub>2</sub> O <sub>2</sub>	282.29	75.0	59.59	5.72	9.88	59.57	5.71	9.92	
30	37		C <sub>14</sub> H <sub>16</sub> Cl <sub>2</sub> N <sub>2</sub> O <sub>2</sub>	315.20	100.0	53.15	4.99	8.86	53.35	5.12	8.88	
31	38		C <sub>14</sub> H <sub>16</sub> F <sub>2</sub> N <sub>2</sub> O <sub>2</sub>	282.29	79.0	59.55	5.73	9.90	59.57	5.71	9.92	
32	39		C <sub>16</sub> H <sub>18</sub> F <sub>3</sub> N <sub>2</sub> O <sub>2</sub>	328.34	oil							o
33	40		C <sub>14</sub> H <sub>16</sub> F <sub>2</sub> N <sub>2</sub> O <sub>2</sub>	282.29	82.5-85	59.72	5.69	9.98	59.57	5.71	9.92	

Exempleno	Compound No.	Structure	Formula	MWt	mp	Cfound	Hfound	Nfound	Cexp	Hexp	Nexp	Note
34	41		C <sub>14</sub> H <sub>16</sub> F <sub>2</sub> N <sub>2</sub> O <sub>2</sub>	282.29	91-92.5	59.58	5.62	9.94	59.51	5.71	9.92	
35	42		C <sub>16</sub> H <sub>18</sub> F <sub>6</sub> N <sub>2</sub> O <sub>2</sub>	382.31	80.5-81.5	50.38	4.25	7.32	50.27	4.22	7.32	
36	43		C <sub>14</sub> H <sub>18</sub> ClN <sub>2</sub> O <sub>3</sub>	288.77	76-78	56.94	6.34	10.25	56.28	6.41	9.37	
37	44		C <sub>14</sub> H <sub>15</sub> ClN <sub>2</sub> O <sub>2</sub>	278.74	123-124	60.88	5.58	9.91	60.33	5.42	10.05	
38	45		C <sub>18</sub> H <sub>24</sub> N <sub>2</sub> O <sub>2</sub>	300.40	94-96	71.89	8.08	9.28	71.97	8.05	9.32	
39	46		C <sub>18</sub> H <sub>28</sub> N <sub>2</sub> O <sub>3</sub>	320.44	oil							b
40	47		C <sub>14</sub> H <sub>19</sub> F <sub>3</sub> N <sub>2</sub> O <sub>3</sub>	282.32	72-73	59.32	6.84	9.81	59.56	6.78	9.92	
41	48		C <sub>18</sub> H <sub>28</sub> N <sub>2</sub> O <sub>2</sub>	302.42	79-80	71.25	8.79	9.36	71.49	8.67	9.26	

Exempleno	Compound No.	Structure	Formula	MWt	mp	Cfound	Hfound	Nfound	Cexp	Hexp	Nexp	Note
42	49		C <sub>14</sub> H <sub>17</sub> F <sub>3</sub> N <sub>2</sub> O <sub>2</sub>	302.30	110.5- 112	55.64	5.77	9.26	55.63	5.67	9.26	
43	50		C <sub>14</sub> H <sub>15</sub> F <sub>3</sub> N <sub>2</sub> O <sub>2</sub>	282.29	94-96	64.29	5.47	10.70	64.11	5.76	10.68	

**Footnotes for Table 1**

Footnote a: IR: 3296, 2980, 2943, 2877, 1638, 1545, 1400, 1377, 1330, 1203, 1166, 1127, 1073, 706  $\text{cm}^{-1}$ .

5 Footnote b: IR 3319, 2963, 2872, 1634, 1549, 1469, 1403, 1327, 1269, 1184, 1130, 1083, 818  $\text{cm}^{-1}$ .

**Example 44. 3-((3-chlorophenyl)methoxy)-azetidine-1-carboxamide (51)**

10 This material was prepared from compound (1) using the procedure described for compound (14) (yield 87%) as a crystalline solid, m.p. 163-165.5°C (diisopropyl ether).

Found: C, 55.49; H, 5.45; N, 11.40.  $\text{C}_{11}\text{H}_{13}\text{ClN}_2\text{O}_2$  requires: C, 54.89; H, 5.44.; N, 11.63.

**Assay Procedures**

15

**Binding to  $\text{CB}_1$  receptors**

The binding of compounds of Formula I to recombinant human  $\text{CB}_1$  receptors was determined *in vitro* by standard methods, with reference to the procedure described by Rinaldi-Carmona *et al.* (Rinaldi-Carmona, M., Pialot, F., Congy, C., Redon, E., Barth, F.,

20 Bachy, A., Breliere, J. C., Soubre, P., LeFur, G., *Life Sci.* 1996, 58(15), 1239-1247).

Membranes were prepared from HEK293 cells expressing recombinant h $\text{CB}_1$  receptors.

Binding assays are performed in a total volume of 250  $\mu\text{L}$ , containing [ $^3\text{H}$ ]-SR-141716A (1 nM final concentration), membranes and test compound. Non-specific binding is determined using CP55,940 (10  $\mu\text{M}$ ). Serial dilutions are performed starting from test compounds as 10

25 mM solutions in DMSO. Compounds are tested over the concentration range  $10^{-10}$  M to  $10^{-5}$  M.  $K_i$  values are calculated from  $\text{IC}_{50}$  values using the Cheng-Prusoff equation.

The thus-determined activity of compounds of formula (I) is shown in Table 2.

30 **Table 2**

Example	$K_i$ (h $\text{CB}_1$ ) nM
Example 6	285

### **Blockade of $\Delta^9$ -THC induced hypolocomotion in mice**

The *in vivo* activity of compounds of formula (1) is assayed for ability to antagonise the reduction in locomotor behaviour induced by acute systemic administration of  $\Delta^9$ -THC in male C57Bl/6 mice. The procedure is as follows.

5

Test compounds are assessed following acute oral or intraperitoneal administration at a dose of 30 mg/kg. Each study utilises a between-subjects design (typically  $n=8$ ) and compares the effects of doses of the test agent to those of vehicle and a positive control.

- 10 The route of test compound administration, drug volume and injection-test-interval are dependent upon the compounds used. 10 min before testing, a 3 mg/kg dose  $\Delta^9$ -THC (or vehicle) is administered to mice by the i.p. route. Automated boxes (AM-1052 activity monitors, Benwick Electronics, Linton Instrumentation) are used to record photocell beam breaks as a measure of locomotor activity. The light beams are arranged on a 7 by 4 matrix
- 15 on a metal grid. 16 grids are connected in series and Perspex boxes, 20 (width) x 40 (length) x 20 (height) cm, with a flat perforated, Perspex lid are placed in each grid. Mice are placed singly in Perspex boxes and the recording of activity in all 16 boxes starts simultaneously. The mice are left undisturbed to explore the novel activity monitor boxes for 15 minutes while beam breaks are recorded.

20

- Locomotor activity data are subjected to one-way analysis of variance (ANOVA) with drug treatment as a between-subjects factor. A significant main effect is followed up by the performance of Dunnett's test in order to assess which treatment mean(s) are significantly different from the control mean. Significant differences between the vehicle /
- 25  $\Delta^9$ -THC group and Test compound /  $\Delta^9$ -THC groups are assessed by Newman-Keuls test. All statistical analyses were performed using Statistica Software, Version 6.0 (Statsoft Inc.) and Microsoft Excel 7.0 (Microsoft Corp.).

### **Regulation of feeding behaviour**


- 30 The *in vivo* activity of compounds of formula (1) is assayed for ability to regulate feeding behaviour by measuring food consumption in male food-deprived Lister-hooded rats as follows.

Test compounds are assessed following acute administration. Each study utilises a between-subjects design (typically  $n=8$ ) and compares the effects of doses of the test agent to those of vehicle and a positive control.

- 5 The anorectic drug sibutramine, or the reference  $CB_1$  receptor antagonist, SR-141716A, normally serves as a positive control. The route of drug administration, drug volume and injection-test-interval are dependent upon the compounds used. The injection-test-interval is the time between dosing and food re-presentation. Typically, animals are fasted such that at the time of food re-presentation food has been withdrawn for an 18-hour period. Food
- 10 consumption is assayed at pre-determined time points (typically 1, 2 and 4 hours after administration). Food intake data are subjected to one-way analysis of variance (ANOVA) with drug as a between-subjects factor. A significant main effect is followed up by the performance of Dunnett's test in order to assess which treatment mean(s) are significantly different from the control mean. All statistical analyses were performed using Statistica
- 15 Software, Version 6.0 (Statsoft Inc.) and Microsoft Excel 7.0 (Microsoft Corp.).

CLAIMS

1. Use of a compound of formula (I)



(I)

wherein:

R<sup>1</sup> is aryl;

R<sup>2</sup> is H, alkyl or aryl; and

R<sup>3</sup> is hydrogen or alkyl;

- or a pharmaceutically acceptable salt or prodrug thereof, in the manufacture of a medicament for the treatment of a disorder mediated by CB<sub>1</sub> receptors.

2. A use according to claim 1 wherein R<sup>1</sup> is a substituted or unsubstituted phenyl or naphthyl.

3. A use according to claim 1 or 2 wherein R<sup>1</sup> has 1, 2 or 3 substituent groups.

4. A use according to claim 1, 2 or 3 wherein R<sup>1</sup> is chlorophenyl or (trifluoromethyl)phenyl.

5. A use according to claim 1, 2, 3 or 4 wherein R<sup>2</sup> is aryl.


6. A use according to any one of claims 1 to 5 wherein R<sup>3</sup> is alkyl.

7. A use according to claim 1 wherein the compound is selected from:  
3-(bis(4-chlorophenyl)methoxy)-N-(2-propenyl)azetidine-1-carboxamide; and  
(R)-3-(bis(4-chlorophenyl)methoxy)-N-(2-hydroxypropyl)azetidine-1-carboxamide.



8. A use according to any preceding claim wherein said medicament comprises a pharmaceutically acceptable carrier and as active ingredient an effective amount of a compound of formula (I).
- 5 9. A method of treatment of a disorder mediated by CB<sub>1</sub> receptors comprising administration to a subject in need of such treatment an effective dose of a compound of formula (I) as defined in any of claims 1 to 8, or a pharmaceutically acceptable salt or prodrug thereof.
- 10 10. A use or method according to any preceding claim wherein the disorder is selected from psychosis, schizophrenia, cognitive disorders, attention deficit disorder, to gastrointestinal disorders, smoking cessation, obesity and other eating disorders associated with excessive food intake, and non-insulin dependant diabetes mellitus.
- 15 11. A use or method according to any of claims 1 to 10 wherein the disorder is obesity.
12. A use or method according to any of claims 1 to 10 for smoking cessation.
13. A use or method according to any of claims 1 to 10 wherein said disorder is a  
20 gastrointestinal disorder.

**PCT/GB2004/001812**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**